

ON THE GROWTH OF ACTIONS OF FREE PRODUCTS

ADRIEN LE BOUDEC, NICOLÁS MATTE BON, AND VILLE SALO

ABSTRACT. If G is a finitely generated group and X a G -set, the growth of the action of G on X is the function that measures the largest cardinality of a ball of radius n in the Schreier graph $\Gamma(G, X)$. In this note we consider the following stability problem: if G, H are finitely generated groups admitting a faithful action of growth bounded above by a function f , does the free product $G * H$ also admit a faithful action of growth bounded above by f ? We show that the answer is positive under additional assumptions, and negative in general. In the negative direction, our counter-examples are obtained with G either the commutator subgroup of the topological full group of a minimal and expansive homeomorphism of the Cantor space; or G a Houghton group. In both cases, the group G admits a faithful action of linear growth, and we show that $G * H$ admits no faithful action of subquadratic growth provided H is non-trivial. In the positive direction, we describe a class of groups that admit actions of linear growth and is closed under free products and exhibit examples within this class, among which the Grigorchuk group.

1. INTRODUCTION

Let G be a finitely generated group, equipped with some finite symmetric generating set S (implicit in what follows). For a G -set X , we denote by $\Gamma(G, X)$ the Schreier graph of the action, with vertex set X and edges (x, sx) for $x \in X$ and $s \in S$. Note that we do not assume the action of G on X to be transitive, so that the graph $\Gamma(G, X)$ need not be connected. Let $\text{vol}_{G,X} : \mathbb{N} \rightarrow \mathbb{N}$ be the function that measures the volume of the largest ball appearing in $\Gamma(G, X)$:

$$\text{vol}_{G,X}(n) = \max_{x \in X} |B_G(n) \cdot x|,$$

where $B_G(n)$ denotes the ball of radius n around the identity in G with respect to the word metric associated to S . For two functions $f_1, f_2 : \mathbb{N} \rightarrow \mathbb{N}$, we write $f_1(n) \leq f_2(n)$ if there exists $C > 0$ such that $f_1(n) \leq C f_2(Cn)$, and $f_1(n) \simeq f_2(n)$ if $f_1(n) \leq f_2(n) \leq f_1(n)$. The function $\text{vol}_{G,X}(n)$ does not depend on the choice of the generating set S up to the equivalence relation \simeq .

Definition 1.1. Given a function $f : \mathbb{N} \rightarrow \mathbb{N}$, we denote by $\mathcal{C}(f)$ the class of finitely generated groups G such that there exists a faithful G -set X such that $\text{vol}_{G,X}(n) \leq f(n)$.

It is not hard to see that the class $\mathcal{C}(f)$ is stable under taking finitely generated subgroups, and under passing to a finite index overgroup [LBMB22a, §1.3]. We use the special notation

Date: January 12, 2024.

Supported by the LABEX MILYON (ANR-10-LABX-0070) of Universite de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007) operated by the French National Research Agency.

$\mathcal{C}(\text{lin})$ for the distinguished case $f(n) \simeq n$, which is the slowest possible growth for a faithful action of an infinite group. The class $\mathcal{C}(\text{lin})$ is richer than it might look at first sight. It is not hard to check that it contains all virtually abelian groups. The fact that non-abelian free groups belong to $\mathcal{C}(\text{lin})$ is well-known and can be traced back to Schreier [Sch27]. A famous example of a group in $\mathcal{C}(\text{lin})$ is the Grigorchuk group (and all Grigorchuk groups G_ω from [Gri84]), see Bartholdi and Grigorchuk [BG00]. Many other interesting groups of dynamical origin are naturally given by an action of linear growth (and hence belong to $\mathcal{C}(\text{lin})$): topological full groups of homeomorphisms of the Cantor set, and Nekrashevych's fragmentation of dihedral groups [Nek18]. More examples of groups in $\mathcal{C}(\text{lin})$ include the lamplighter group $C_p \wr \mathbb{Z}$, the Houghton groups, or Neumann's groups from [Neu37].

In [Sal21] the third author investigated the class of subgroups of the topological full group of full shifts over \mathbb{Z} (which is contained in $\mathcal{C}(\text{lin})$), and gave sufficient conditions under which graph products of such groups remain inside that class [Sal21, Theorem 3]. That result implies in particular that the class $\mathcal{C}(\text{lin})$ contains all right angled Artin groups [Sal21, Theorem 1], and hence all their subgroups as well. Combined with results of many authors, this is a vast source of examples of groups in $\mathcal{C}(\text{lin})$, as many groups are known to embed (upon to passing to a finite index subgroup) in a right angled Artin groups. For example, this is the case for instance surface groups, and more generally any hyperbolic group acting geometrically on a CAT(0) cube complex, as follows from Agol's final step in the solution of the virtual Haken conjecture [Ago13] combined with the previous results of Haglund and Wise [HW08].

Motivated by this last class of examples, we consider here the question whether the class $\mathcal{C}(\text{lin})$ (or the classes $\mathcal{C}(f)$ for more general f) is closed under free products, or more generally graph products. There is a natural strategy to attempt to answer that question affirmatively. Given two groups G_1 and G_2 and X_i a G_i -set, $i = 1, 2$, one may consider any set $X = X_1 \sqcup_{A_1 \cong_{\phi} A_2} X_2$ obtained by identifying suitable subsets $A_1 \subset X_1$ and $A_2 \subset X_2$ via a bijection $\phi: A_1 \rightarrow A_2$. The action of each G_i extends to X trivially outside of the copy of X_i in X . This defines an action of the free product $G = G_1 * G_2$. On the one hand, if the subsets along which the X_i are glued are chosen wisely, one can hope to control the growth function $\text{vol}_{G,X}(n)$ in terms of $\text{vol}_{G_i, X_i}(n)$, $i = 1, 2$. On the other hand, if the gluing is sufficiently generic, the action of G on X is likely to be faithful. The question therefore becomes whether, for G_1, G_2 in $\mathcal{C}(f)$, the tension between these two conditions can be conciliated. A naive implementation of this strategy (suitably adapted to more general graph products) gives the following. We refer to §2.1 for the definition of graph products.

Proposition 1 (Proposition 2.7). *Let $G_1, \dots, G_k \in \mathcal{C}(f)$. Then the free product $G = G_1 * \dots * G_k$ belongs to $\mathcal{C}(g)$, where $g(n) = nf(n)$. More generally, any graph product of the G_i 's belongs to $\mathcal{C}(g)$.*

Proposition 1 is very often not optimal. In many cases, appropriate choices allow to show that a graph product of groups in $\mathcal{C}(f)$ remains in $\mathcal{C}(f)$. In §2.3 we discuss a sufficient condition for this to be the case. That condition is an elaboration of a condition considered in [Sal21]. Proposition 2.13 provides in particular a direct and elementary proof of the fact

that right-angled Artin groups belong to $\mathcal{C}(\text{lin})$ (which essentially follows the same lines as the proof in [Sal21], but avoids the technicalities coming from the full group setting). More generally, Proposition 2.13 allows to exhibit examples of graph products in the class $\mathcal{C}(\text{lin})$:

Proposition 2. *The free product $G_1 * \cdots * G_k$ belongs to $\mathcal{C}(\text{lin})$ whenever each free factor G_i is isomorphic to one of the following:*

- a finite group;
- a finitely generated abelian group;
- a finitely generated subgroup of a right-angled Artin group;
- the Grigorchuk group;
- the lamplighter group $(\mathbb{Z}/p\mathbb{Z}) \wr \mathbb{Z}$ for $p \geq 2$.

More generally, the graph product of any finite family of groups in the list above belongs to $\mathcal{C}(\text{lin})$.

1.1. The main results. The main goal of this note is to provide examples that show that the bound obtained from the simple construction from Proposition 1 is sharp in general, even for free products. We consider two families of examples.

The first are topological full groups of minimal group actions on the Cantor set, more precisely their alternating subgroups in the sense of Nekrashevych [Nek19] (see §3.4 for definitions). Following [LBMB22a], we shall say that a finitely generated group G has a **Schreier growth gap** $f(n)$ if every faithful G -set X satisfies $\text{vol}_{G,X}(n) \geq f(n)$.

Theorem 1. *Let $\mathcal{G} \curvearrowright \mathcal{X}$ be a minimal expansive action of a finitely generated group on the Cantor set, and set $f(n) = \text{vol}_{\mathcal{G},\mathcal{X}}(n)$. Let G be the alternating full group of the action (so that G is finitely generated and belongs to $\mathcal{C}(f(n))$, see §3.4). Then for every non-trivial finitely generated group H , the group $G * H$ has a Schreier growth gap $nf(n)$.*

A relevant special case in the previous theorem is $\mathcal{G} = \mathbb{Z}$ (in that case G coincides with the commutator subgroup of the topological full group, by the results of Matui [Mat06]). In that case we obtain examples of groups $G \in \mathcal{C}(\text{lin})$ such that $G * H$ has a Schreier growth gap n^2 for any non-trivial group H . The quadratic gap is optimal in general, as Proposition 1 says that $G * H$ has a faithful action of growth $\simeq n^2$ for instance when H is cyclic and non-trivial.

Our second family of examples are the Houghton groups $H_r, r \geq 2$. The group H_r is finitely generated and belongs to $\mathcal{C}(\text{lin})$ for every $r \geq 2$ (see §3.3).

Theorem 2. *Let $G = H_r$ be the Houghton group on $r \geq 2$ rays. Then for every non-trivial finitely generated group H , the group $G * H$ has a Schreier growth gap n^2 .*

1.2. Outline of the proof of Theorems 1 and 2. . These two results share the same proof mechanism. It relies on the description of the confined subgroups of the groups G in their statements, obtained in previous works of the first two authors. Recall that a subgroup H of a group G is confined if the set of conjugates of H does not accumulate on the trivial subgroup in the space $\text{Sub}(G)$ of subgroups of G , endowed with the Chabauty topology. When G is finitely generated, convergence in the space $\text{Sub}(G)$ can be interpreted in the space of marked Schreier graphs on the corresponding coset spaces G/H . The study

of confined subgroups of a group G leads (among other applications) to results on the growth function $\text{vol}_{G,X}(n)$ of G -actions, such as the existence of a Schreier growth gap [MB18, LBMB23, LBMB22a]. For G as in Theorem 1 or 2, a classification of the confined subgroups of G has been obtained respectively in [MB18] and [LBMB22a]. Using these results, we show that every faithful action of G whose growth is close to the growth of the natural defining action of G (respectively on the Cantor set or on the bouquet of rays), must be “almost” conjugate to it. A common feature of the two situations is that the group G contains elements whose support is a very sparse subset of the Schreier graph of the natural action. We exploit this fact and the previous result about actions of small growth of G to show that in the graph for any faithful action of $G * H$, there must be short jumps between regions that are far in the graph of the restricted action of G . This is that phenomenon that forces an additional factor n in the growth. The detailed proofs are given in Section 3.

It is worth comparing the case of the Grigorchuk group in Proposition 2 with Theorem 1. The Grigorchuk group and topological full groups share various common features: in particular they appear through a micro-supported action on a compact space, a condition which plays an important role in the study of confined subgroups (indeed the confined subgroups of the Grigorchuk group are also understood [LBMB23]). Also Theorem 1 is applicable to many fragmentations of dihedral groups in the sense of Nekrashevych [Nek18] (a family of groups to which the Grigorchuk group also belongs). Despite these similarities, here they exhibit an opposite behaviour. The main reason for this difference is the fact that its action on the vertices of the binary tree satisfies a condition that we call large displacement (Definition 2.9), which means that every non-trivial element of the group must move some vertex of the tree by a distance (on the Schreier graph of the corresponding level) comparable to the total diameter of the Schreier graph (Proposition 2.14). Although the proof of this fact is not difficult, it relies on rather specific features of the Grigorchuk group.

2. STABILITY RESULTS

2.1. Graph products. Let I be a set, and let Δ denote the diagonal in I^2 . Let $c : I^2 \setminus \Delta \rightarrow \{0, 1\}$ such that $c(i, j) = c(j, i)$ for all $i \neq j$. If $(G_i)_{i \in I}$ a family of groups indexed by I , the **graph product** $P = (G_i)_{i \in I}^c$ of the family $(G_i)_{i \in I}$ associated to c is the quotient of the free product $*_I G_i$ by the relations $[G_i, G_j]$ for all i, j such that $c(i, j) = 1$. We say that P is a **finite graph product** if I is finite.

2.2. Preliminaries.

Definition 2.1. Let $q \geq 1$. Let $(X_1, \dots, X_q) = (X_k)_{k \leq q}$ be a q -tuple of graphs, and for every $k \leq q$ let (e_k, s_k) be two distinct points of X_k . The **gluing** of the q -tuple $(X_k, e_k, s_k)_{k \leq q}$ is the graph obtained by taking the disjoint union $X_1 \sqcup \dots \sqcup X_q$, and identifying s_k and e_{k+1} for every $1 \leq k \leq q - 1$. It is denoted $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$.

(The points e_1 and s_q do not play any role, but we keep two based points in X_1 and X_q as well in order to simplify notation.) Note that for every $k \leq q$ the map from X_k to $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ is a graph isomorphism onto its image. In the sequel we identify X_k

with its image in $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$. Note also that the condition $e_k \neq s_k$ ensures that the images of X_k and X_{k+r} in $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ are disjoint for all $r \geq 2$.

Lemma 2.2. *Let $(X_k)_{k \leq q}$ be a q -tuple of graphs, and suppose that every ball of radius $n \geq 1$ in X_k has cardinality at most $f(n)$ for every $k \leq q$. Then every ball of radius n in $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ has cardinality at most $(2n + 1) \cdot f(n)$.*

Proof. Every ball B of radius n in $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ intersects at most $2n + 1$ members of $(X_k)_{k \leq q}$, and is covered by $2n + 1$ balls of radius at most n within these X_k . The statement follows. \square

In the sequel we fix a set I , a function $c : I^2 \setminus \Delta \rightarrow \{0, 1\}$ such that $c(i, j) = c(j, i)$ for all $i \neq j$, and a family of graphs \mathcal{F} such that \mathcal{F} admits a partition indexed by I , with blocks denoted \mathcal{F}_i , $i \in I$. For $X \in \mathcal{F}$, we write $i(X)$ for the unique element of I such that $X \in \mathcal{F}_{i(X)}$.

Definition 2.3. A q -tuple $(X_k)_{k \leq q}$ of elements of \mathcal{F} is **c -admissible** if $i(X_k) \neq i(X_{k+1})$ and $c(i(X_k), i(X_{k+1})) = 0$ for every $k = 1, \dots, q - 1$. By extension we also say that $(X_k, e_k, s_k)_{k \leq q}$ is c -admissible if $(X_k)_{k \leq q}$ is c -admissible.

Definition 2.4. We let $\mathcal{G}(\mathcal{F}, c)$ be the disjoint union of all $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$, where $q \geq 1$ is any positive integer, $(X_k)_{k \leq q}$ is any c -admissible q -tuple, and $(e_k, s_k)_{k \leq q}$ is any sequence such that e_k, s_k are distinct elements of X_k for every $k \leq q$.

Now suppose that $(G_i)_{i \in I}$ is a family of groups, and for every $i \in I$ there is an action of G_i on X for every $X \in \mathcal{F}_i$. Let $P = (G_i)_{i \in I}^c$ be the graph product of $(G_i)_I$ associated to c . For every c -admissible $(X_k, e_k, s_k)_{k \leq q}$, there is a natural G_i -action on $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ that extends the G_i -action on each X_k such that $i(X_k) = i$, that is defined by declaring that G_i acts trivially on $X_k \setminus \{e_k, s_k\}$ for every k such that $i(X_k) \neq i$. This is well defined because of the property that no two consecutive X_k can be in \mathcal{F}_i , guaranteed by the assumption $i(X_k) \neq i(X_{k+1})$ in Definition 2.3. This defines an action of the free product $*_I G_i$ on $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$. The fact that $(X_k, e_k, s_k)_{k \leq q}$ is c -admissible ensures that if $c(i, j) = 1$, then the groups G_i and G_j act on $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ with disjoint support. Hence the action of $*_I G_i$ on $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ factors through the graph product $P = (G_i)_{i \in I}^c$.

Remark 2.5. Suppose for every $i \in I$ the group G_i is generated by a subset S_i , and that the graph structure on each $X \in \mathcal{F}_i$ is the Schreier graph of the G_i -action on X . Then the graph $\mathcal{G}((X_k, e_k, s_k))$ is essentially the Schreier graph of the action of $P = (G_i)_{i \in I}^c$ associated to the generating subset $\cup_I S_i$. The only difference is that loops should be added at the places where the action of the G_i 's has been extended in a trivial way.

Lemma 2.6. *Suppose that for every $i \in I$, the G_i -action on $\bigsqcup_{X \in \mathcal{F}_i} X$ is faithful. Then the action of the graph product $P = (G_i)_{i \in I}^c$ on $\mathcal{G}(\mathcal{F}, c)$ is faithful.*

Proof. Let g be a non-trivial element of P . Consider the decompositions of the form

$$g = g_n \cdots g_1,$$

where for each s we have that g_s is non-trivial and there is $i_s \in I$ (necessarily unique) such that $g_s \in G_{i_s}$. Among all such decomposition, we choose one such that n is minimal. Since g is non-trivial, $n \geq 1$. We define a sequence (r_1, \dots, r_q) inductively by setting $r_1 = 1$, and defining r_{k+1} as the smallest $s \geq r_k + 1$ such that $c(i(s), i(r_k)) = 0$ (equivalently, $G_{i(s)}$ does not commute with $G_{i(r_k)}$). We also define $r_{q+1} = n + 1$. Minimality of n implies that for every $1 \leq k \leq q$ and every $r_k + 1 \leq s \leq r_{k+1} - 1$, we have $g_s \notin G_{i(r_k)}$. Let $\gamma_k = g_{r_{k+1}-1} \cdots g_{r_k}$ for $1 \leq k \leq q$, so that $g = \gamma_q \cdots \gamma_1$.

Since G_i acts faithfully on $\bigsqcup_{X \in \mathcal{F}_i} X$ for every $i \in I$, for every $k \leq q$ one can find $X_k \in \mathcal{F}_{i_{r_k}}$ and $x_k \in X_k$ such that $g_{r_k}(x_k) \neq x_k$. We set $e_k = x_k$, $s_k = g_{r_k}(x_k)$. We look at how the element g acts on $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$. Since g_s does not belong to $G_{i(r_k)}$ for every $r_k + 1 \leq s \leq r_{k+1} - 1$, g_s acts trivially on s_k . Hence $\gamma_k(e_k) = s_k$ for every $k \leq q$, and hence it follows that $g(e_1) = s_q$. So if $q = 1$ then g acts non-trivially since $e_1 \neq s_1$. If $q \geq 2$ then X_q and X_1 are either disjoint (when $q \geq 3$), or intersect only along $s_1 = e_2$ if $q = 2$. Hence in all cases the element g moves e_1 . So for every non-trivial element g of P , there exists a $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ on which g acts non-trivially. So P acts faithfully on $\mathcal{G}(\mathcal{F}, c)$. \square

Recall that $\mathcal{C}(f)$ has been defined in the introduction (Definition 1.1).

Proposition 2.7. *Let $f: \mathbb{N} \rightarrow \mathbb{N}$, and let $(G_i)_I$ be a finite collection of groups in the class $\mathcal{C}(f)$. Then any graph product $P = (G_i)_{i \in I}^c$ belongs to $\mathcal{C}(g)$ with $g(n) = nf(n)$.*

Proof. For each $i \in I$ there is a G_i -set X_i such that the G_i -action on X_i is faithful and verifies $\text{vol}_{G_i, X_i}(n) \leq f(n)$. The statement then follows from Lemma 2.2 and Lemma 2.6 (and Remark 2.5), applied to $\mathcal{F}_i = \{X_i\}$ and $\mathcal{F} = \bigsqcup_I \mathcal{F}_I$. \square

2.3. A subclass closed under graph products. In this section we prove that, under a mild assumption on the function f , a certain subclass of $\mathcal{C}(f)$ is closed under finite graph products (Proposition 2.13). The definition of this subclass is inspired by the work of the third author [Sal21], and Proposition 2.13 elaborates on [Sal21, Theorem 3]. In particular Proposition 2.13 provides an elementary proof of the result from [Sal21] that finitely generated right-angled Artin groups admit a faithful action with linear growth. (Note that Theorem 3 in [Sal21] is stronger than that).

For an increasing function $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$, consider the following condition:

$$(1) \quad \exists C_1 > 0; \forall k \geq 1 \forall \rho_1, \dots, \rho_k \geq 0, \sum_{i \leq k} f(\rho_i) \leq C_1 f\left(C_1 \sum_{i \leq k} \rho_i\right).$$

It is a rather mild condition. A sufficient condition for (1) to hold is that $n \mapsto f(n)/n$ is increasing. For instance $f(n) = n^\alpha$ satisfies (1) for every $\alpha \geq 1$.

Lemma 2.8. *Let $(X_k)_{k \leq q}$ be a q -tuple of finite graphs, and e_k, s_k distinct elements of X_k for every $k \leq q$. Suppose that there is an increasing function $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ and $C > 0$ such that:*

- (1) every ball of radius n in X_k has cardinality at most $Cf(Cn)$ for every $k \leq q$.
- (2) f satisfies (1).
- (3) $\text{diam}(X_k) \leq Cd(e_k, s_k)$ for every $k \leq q$.

Then there is a constant $C' > 0$ depending only on C and f such that every ball of radius n in $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$ has cardinality at most $C'f(C'n)$.

Proof. Let C_1 coming from the fact that f satisfies (1). Let $n \geq 1$. Let x be a vertex in $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$, and let ℓ such that x belongs to X_ℓ . Let $B(n)$ denote the ball of radius n around x in $\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$. Let $X_{\ell-\ell_1}, \dots, X_\ell, \dots, X_{\ell+\ell_2}$ be the members of $(X_k)_{k \leq q}$ that $B(n)$ intersects. Certainly we have

$$d(e_{\ell+1}, s_{\ell+1}) + \dots + d(e_{\ell+\ell_2-1}, s_{\ell+\ell_2-1}) \leq n$$

and

$$d(e_{\ell-1}, s_{\ell-1}) + \dots + d(e_{\ell-\ell_1+1}, s_{\ell-\ell_1+1}) \leq n,$$

and $B(n)$ is contained in the union of $X_{\ell-\ell_1+1}, \dots, X_{\ell-1}, X_{\ell+1}, \dots, X_{\ell+\ell_2-1}$ and the intersection between $B(n)$ and $X_\ell \cup X_{\ell-\ell_1} \cup X_{\ell+\ell_2}$:

$$|B(n)| \leq \sum_{k=1}^{\ell_1-1} |X_{\ell-k}| + \sum_{k=1}^{\ell_2-1} |X_{\ell+k}| + |B_{X_\ell}(x, n)| + |B_{X_{\ell-\ell_1}}(s_{\ell-\ell_1}, n)| + |B_{X_{\ell+\ell_2}}(e_{\ell+\ell_2}, n)|.$$

Conditions 1 and 3 ensure $|X_k| \leq Cf(C \text{diam}(X_k)) \leq Cf(C^2 d(e_k, s_k))$ for every $k \leq q$. Hence we obtain

$$\begin{aligned} |B(n)| &\leq C \sum_{k=1}^{\ell_1-1} f(C^2 d(e_{\ell-k}, s_{\ell-k})) + C \sum_{k=1}^{\ell_2-1} f(C^2 d(e_{\ell+k}, s_{\ell+k})) + 3Cf(Cn) \\ &\leq C_1 Cf \left(C_1 C^2 \sum_{k \leq \ell_1-1} d(e_{\ell-k}, s_{\ell-k}) \right) + C_1 Cf \left(C_1 C^2 \sum_{k \leq \ell_2-1} d(e_{\ell+k}, s_{\ell+k}) \right) + 3Cf(Cn) \\ &\leq 2C_1 Cf(C_1 C^2 n) + 3Cf(Cn), \end{aligned}$$

where in the second inequality we have used that f satisfies (1), and the third inequality follows from the above upper bounds $\sum d(e_{\ell-k}, s_{\ell-k}), \sum d(e_{\ell+k}, s_{\ell+k}) \leq n$. \square

Definition 2.9. Let $G = \langle S \rangle$ be a finitely generated group, and X a G -set. We say that the G -action on X has **large displacement** if all G -orbits in X are finite, and if there is $C > 0$ such that for every non-trivial element $g \in G$, there is a G -orbit O in X and $x \in O$ such that $gx \neq x$ and $\text{diam}(\Gamma(G, O)) \leq Cd(x, gx)$.

Here by $\Gamma(G, O)$ we mean the Schreier graph of the G -action on O (which is a connected component in the larger graph $\Gamma(G, X)$). It is easily checked that this condition does not depend on the choice of S (although C can depend on S). Note that the definition forces in particular the group G to be residually finite.

Definition 2.10. We denote by $\mathcal{C}_{LD}(f) \subset \mathcal{C}(f)$ the class of groups admitting an action with large displacement and of growth at most $f(n)$.

Example 2.11.

- (1) Every faithful action of a finite group has large displacement.

- (2) The action of \mathbb{Z} on $\bigsqcup_{n \geq 1} \mathbb{Z}/n\mathbb{Z}$ has large displacement. Hence \mathbb{Z} belongs to $\mathcal{C}_{LD}(\text{lin})$.
- (3) If G_1, \dots, G_k admit actions with large displacement respectively on X_1, \dots, X_k , then the action of $G_1 \times \dots \times G_k$ on $X_1 \sqcup \dots \sqcup X_k$ has large displacement. Hence each class $\mathcal{C}_{LD}(f)$ is closed under direct products.

Here is a slightly more elaborated example:

Proposition 2.12. *For every $d \geq 1$ and $p \geq 2$, the wreath product $G = (\mathbb{Z}/p\mathbb{Z}) \wr \mathbb{Z}^d$ belongs to $\mathcal{C}_{LD}(f)$ for $f(n) = n^d$.*

Proof. We denote elements of G as pairs (f, u) , with $f: \mathbb{Z}^d \rightarrow \mathbb{Z}/p\mathbb{Z}$ of finite support and $u \in \mathbb{Z}^d$. For $m \geq 2$, set $H_m = \{(f, u): \sum_{v \in (m\mathbb{Z})^d} f(v) = 0, u \in m\mathbb{Z}^d\}$. Then H_m is a finite index subgroup of G , of index pm . Set $X_m := G/H_m$ and $X = \sqcup_m X_m$. We check that the action of G on X has large displacement and satisfies $\text{vol}_{G,X}(n) \simeq n^d$. Consider the standard generating set $S = \{(\delta_0, 0), (0, e_i), i = 1, \dots, d\}$, where $\delta_0: \mathbb{Z}^d \rightarrow \mathbb{Z}/p\mathbb{Z}$ takes the value 1 on 0 and 0 elsewhere, and e_1, \dots, e_d is the standard basis of \mathbb{Z}^d . Note first that the map $G \rightarrow (\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/m\mathbb{Z})^d$ that sends $(f, u) \in G$ to $(\sum_{v \in m\mathbb{Z}^d} f(v), u \bmod m)$ descends to a bijection of the coset space G/H_m with $(\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/m\mathbb{Z})^d$, so that we may identify X_m with $(\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/m\mathbb{Z})^d$. Under this identification, the action of G on X_m coincides with the standard wreath product action of the natural quotient $(\mathbb{Z}/p\mathbb{Z}) \wr (\mathbb{Z}/m\mathbb{Z})^d$. Explicitly, the action of elements in the generating set S is described as follows: the lamp generator $(\delta_0, 0)$ permutes cyclically $(\mathbb{Z}/p\mathbb{Z}) \times \{0\}$ and acts trivially elsewhere; each element $(0, e_i)$ maps $(r, u \bmod m)$ to $(r, u + e_i \bmod m)$. In particular the Schreier graph of the G -action on X_m is isomorphic (ignoring loops) to the graph obtained taking a cycle of length p and gluing to each point a copy of the standard Cayley graph of $(\mathbb{Z}/m\mathbb{Z})^d$. It follows that $\text{vol}_{G,X}(n) \simeq n^d$, and that $C_1 m \leq \text{diam}(X_m) \leq C_2 m$ for some constants C_1, C_2 . For the large displacement property, let $g = (f, v)$ be a non-trivial element of G . Suppose first that $v \neq 0$, and choose $m = 2|v|$, where $|\cdot|$ denotes the standard word metric of \mathbb{Z}^d . Then, since the natural projection of X_m to $(\mathbb{Z}/m\mathbb{Z})^d$ is equivariant, for every $x \in X_m$ we have $d(x, gx) \geq |v| \geq \frac{1}{2} C_1 \text{diam}(X_m)$. Suppose now that $v = 0$. Let u be such that $f(u) \neq 0$ and with $|u|$ maximal, and choose again $m > 2|u|$. Let $x \in X_m \simeq (\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/m\mathbb{Z})^d$ be the point $x = (0, u \bmod m)$. Then $gx = (f(u), u \bmod m)$. From the description of the graph $\Gamma(G, X_m)$ one can see that $d(x, gx) \geq 2|u| \geq C_1 \text{diam}(X_m)$. \square

Our main motivation for considering the class $\mathcal{C}_{LD}(f)$ is the following:

Proposition 2.13. *If $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is increasing and satisfies (1), then the class $\mathcal{C}_{LD}(f)$ is closed under finite graph products.*

Proof. Let $(G_i)_I$ be a finite collection of groups in $\mathcal{C}_{LD}(f)$. For every $i \in I$ let Y_i be a G_i -set such that the G_i -action on Y_i has large displacement and growth at most $f(n)$. Let S_i be a finite generating subset of G_i and C_i a constant as in Definition 2.9. Let $P = (G_i)_{i \in I}^c$ be a graph product, generated by $S = \cup S_i$. Set $C = \max C_i$.

For $i \in I$, let \mathcal{F}_i be the set of G_i -orbits in Y_i , and \mathcal{F} the disjoint union of the \mathcal{F}_i . We consider the subset of $\mathcal{G}(\mathcal{F}, c)$, denoted by Y , consisting of the disjoint union of all

$\mathcal{G}((X_k, e_k, s_k)_{k \leq q})$, where $q \geq 1$, $(X_k)_{k \leq q}$ is a c -admissible q -tuple of elements of \mathcal{F} , and e_k, s_k are distinct elements of X_k with the condition that $\text{diam}(X_k) \leq Cd(e_k, s_k)$ for every $k \leq q$. The group P acts on Y , and every P -orbit in Y is finite because all the members of \mathcal{F} are finite. Moreover Lemma 2.8 ensures $\text{vol}_{P,Y}(n) \leq f(n)$. Hence to conclude the proof it suffices to see that the P -action on Y has large displacement.

Let g be a non-trivial element of P . We repeat verbatim the argument in the proof of Lemma 2.6. We find a decomposition $g = \gamma_q \cdots \gamma_1$ and a c -admissible $(X_k, e_k, s_k)_{k \leq q}$ such that $\gamma_k(e_k) = s_k$ for every $k \leq q$ (so that $g(e_1) = s_q$). The assumption that the G_i -action on Y_i has large displacement allows to ensure that $\text{diam}(X_k) \leq Cd(e_k, s_k)$ for every $k \leq q$. Hence if we write $X = \mathcal{G}((X_k, e_k, s_k)_{k \leq q})$, then $X \in Y$ and

$$\text{diam}(X) \leq \sum_{k=1}^q \text{diam}(X_k) \leq C \sum_{k=1}^q d(e_k, s_k) = Cd(e_1, s_q).$$

□

2.4. The Grigorchuk group. In this section we prove that the Grigorchuk group belongs to $\mathcal{C}_{LD}(\text{lin})$. We denote by $T = \{0, 1\}^*$ the rooted tree of finite binary words, where each word w is connected by an edge to $wx, x \in \{0, 1\}$. Recall that the Grigorchuk group G is the subgroup of the automorphism group $\text{Aut}(T)$ generated by the set of automorphisms $S = \{a, b, c, d\}$ given by the recursive rules

$$\begin{aligned} a(0w) &= 1w, & a(1w) &= 0w; \\ b(0w) &= 0a(w), & b(1w) &= 0c(w); \\ c(0w) &= 0a(w), & c(1w) &= 1d(w); \\ d(0w) &= 0w, & d(1w) &= 1b(w). \end{aligned}$$

We denote by Γ_n the Schreier graph of the action of G on the level $\{0, 1\}^n$ of the tree. We recall basic properties of the structure of these graphs, see [BG00] for a more detailed description. Each Γ_n is isometric to an interval of length $2^n - 1$ in \mathbb{Z} (with loops and multiple edges, which won't be important for us). Along these segments, binary words are ordered following the *Gray code* ordering, namely each $w \in \{0, 1\}^n$ is connected by an edge in Γ_n to the word obtained by changing its first digit, and to the word obtained by changing the digit that follows the first appearance of 0 in w (if the latter exists). Thus all words have exactly two neighbours, with the exception of the words $11 \cdots 11$ and $11 \cdots 10$, which lie at the extreme points of Γ_n (we shall picture this with $11 \cdots 11$ as the leftmost point and $11 \cdots 10$ as the rightmost point). Let us denote by d_n the associated distance on $\{0, 1\}^n$. The map $p_n: \{0, 1\}^n \rightarrow \{0, 1\}^{n-1}$ that erases the last digit induces a covering map $p_n: \Gamma_n \rightarrow \Gamma_{n-1}$, which corresponds to folding Γ_n around its middle edge.

Proposition 2.14. *Let G be the Grigorchuk group and retain the above notation. Then for every non-trivial element $g \in G$, there exists $n \geq 1$ and $w \in \{0, 1\}^n$ such that $d_n(gw, w) \geq \frac{1}{8} \text{diam}(\Gamma_n)$. In particular G belongs to $\mathcal{C}_{LD}(\text{lin})$.*

Recall that **portrait** of an element $g \in G$ is the collection permutations $(\sigma_w)_{w \in T}$, where each $\sigma_w \in \text{Sym}(2)$ specifies the action of g on the two children of w . In formulas, we have

$g(wx) = g(w)\sigma_w(x)$ for $x \in \{0, 1\}$. We say that w is an **active vertex** of g if σ_w is non-trivial. We will rely on a result in [AS07], where Arzhantseva and Šunić provide an explicit list of all the restrictions of portraits of elements of Grigorchuk groups to finite subtrees of depth 3 (and show that this characterises the closure of G in $\text{Aut}(T)$).

Proof of Proposition 2.14. Let $g \in G$ be non-trivial. Let m be the smallest level containing active vertices. We can suppose that G fixes $\{0, 1\}^n$ for all $n \leq 3$, or the conclusion holds true trivially for g , since $\text{diam}(\Gamma_n) \leq 7$ for $n \leq 3$ and any point which is not fixed by g is moved at distance at least 1. Hence $m \geq 3$. Choose $w \in \{0, 1\}^m$ active, let w_0 be the projection of w at level $m - 3$, and T_0 be the finite rooted at w_0 and containing all its descendants up to 3 levels. Hence w is a leaf of T_0 , and all active vertices for g on T_0 must be leaves, by minimality of m . It follows from Theorem 1 in [AS07] that at least another leaf of T_0 , distinct from w , must be active. It follows that we can find two distinct active vertices $w, v \in \{0, 1\}^m$ having the same projection at level $m - 3$. Consider now the action on level $m + 1$. Since g fixes level m , it must preserve all fibers of the covering map $p_{m+1}: \Gamma_{m+1} \rightarrow \Gamma_m$, which consist each of two points, and it acts non-trivially on each of the two fibers $p_{m+1}^{-1}(w), p_{m+1}^{-1}(v)$. Let $I \subset \Gamma_m$ be the interval ending at the right-most points of length $2^{m-3} \geq \frac{1}{8} \text{diam}(\Gamma_m)$. For any vertex $u \notin I$, the two points in $p_{m+1}^{-1}(u)$ are at distance at least $2|I| \geq \frac{1}{8} \text{diam}(\Gamma_{m+1})$ in Γ_{m+1} . On the other hand the projection map from $\Gamma_m \rightarrow \Gamma_{m-3}$ is injective on I , and since w, v have the same image, they can't both belong to I . Hence the desired conclusion follows, for $n = m + 1$. \square

2.5. The proof of Proposition 2. Note that Proposition 2 from the introduction follows from Proposition 2.13, together with the combination of Example 2.11, Proposition 2.12 and Proposition 2.14.

3. SCHREIER GROWTH GAPS FOR FREE PRODUCTS

3.1. A criterion. Let G be a finitely generated group endowed with a finite symmetric generating set S . If X is a G -set, we endow X with the its Schreier graph structure. The associated simplicial distance is denoted $d_{G,X}$. We denote by $B_{G,X}(x, n)$ the corresponding balls. The **support** of an element $g \in G$ in X is the set $\text{supp}_X(g) = \{x \in X : gx \neq x\}$. Given $R > 0$, we shall say that two subsets $A, B \subset X$ are R -separated if $d_{G,X}(a, b) \geq R$ for every $a \in A$ and $b \in B$. Finally the **R -coarse connected components** of a subset $A \subset X$ are the equivalence classes of the equivalence relation on A generated by the pairs (a_1, a_2) such that $d_{G,X}(a_1, a_2) \leq R$.

Definition 3.1. Let G be a finitely generated group endowed with a finite symmetric generating set S . Let $\alpha, \beta: \mathbb{N} \rightarrow \mathbb{N}$ be functions such that $\alpha(n) \leq \beta(n)$. We say that G satisfies the sparse support condition at scale (α, β) , if for every faithful G -set X such that $\text{vol}_{G,X}(n) \not\asymp \beta(n)$, there exists constants $C, D > 0$ such that for every $R > 0$, there exists a non-trivial element $g \in G$ satisfying the following

- (C1) We have $d_{G,X}(x, gx) \leq D$ for every $x \in X$.
- (C2) Every D -coarse connected component of $\text{supp}_X(g)$ has diameter at most C .

- (C3) Any two distinct D -coarse connected components of $\text{supp}_X(g)$ are R -separated.
(C4) For every $x \in \text{supp}_X(g)$, we have $|B_{G,X}(x, R)| \geq \frac{1}{C} \alpha(\frac{1}{C}R)$.

This condition does not depend on the choice of a finite generating set S for G . Note that condition (C4) implies in particular that G satisfies a Schreier growth gap $\alpha(n)$. The sparse support condition implies that this gap can be improved for any non-trivial free product with G , as follows.

Proposition 3.2. *Suppose that G satisfies the sparse support condition at scale (α, β) . Then for every non-trivial finitely generated group H , the group $G * H$ has a Schreier growth gap $\min(n\alpha(n), \beta(n))$.*

Proof. Set $L = G * H$. We fix a finite generating set of L of the form $S \cup T$, where S is a generating set of G and H is a generating set of H . Let $t \in T$ be a non-trivial. Let X be a faithful L -set. We shall consider on X the Schreier graph distance by $d_{L,X}$, as well as the distance $d_{G,X}$ induced by restricting the action to G . Suppose that $\text{vol}_{G,X}(n) \not\geq \beta(n)$ (else, the desired conclusion is true trivially). Let C, D be as in Definition 3.1, fix $R > 0$ (which we may assume is even), and let $g \in G$ be the corresponding element. Consider the commutator $h = [g, t]$. Note that $\text{supp}_X(h)$ is contained in the 1-neighbourhood of $\text{supp}_X(g)$ (with respect to the distance $d_{L,X}$).

Since h has infinite order and the action is faithful, we can find $x_0 \in X$ such that the points $x_n = h^n(x_0)$ are pairwise distinct for $n = 0, \dots, R$. Note that $d_{L,X}(x_n, x_{n+1}) \leq 2D + 2$, by condition (C1). For each point x_n we choose $y_n \in \text{supp}_X(g)$ such that $d_{L,X}(x_n, y_n) \leq 1$. Then all points y_0, \dots, y_R belong to the ball $B_{L,X}(x_0, D_1R)$, with $D_1 = 2D + 3$. Since all D -coarse connected components of $\text{supp}_X(g)$ with respect the distance $d_{G,X}$ have diameter bounded by C , their 1-neighbourhood in the distance $d_{L,X}$ has cardinality uniformly bounded by some constant $C_1 > 0$ (not depending on R).

Now from the fact that the points x_n are all distinct, we deduce that at most C_1 points y_0, \dots, y_R belong to the same D -coarse connected component. It follows that from $\{y_0, \dots, y_n\}$ we can extract a collection of points $z_1, \dots, z_{\lfloor R/C_1 \rfloor}$ belonging to distinct D -coarse connected components. Then the balls $B_{G,X}(z_i, R/2)$ are pairwise disjoint by (C3), moreover they are contained in $B_{L,X}(x_0, D_2R)$, with $D_2 = D_1 + 1/2$. From (C4) we deduce that

$$|B_{L,X}(x_0, D_2R)| \geq \sum_{i=1}^{\lfloor R/C_1 \rfloor} |B_{G,X}(z_i, R/2)| \geq \frac{1}{C_2} R \alpha\left(\frac{1}{C_2}R\right),$$

for some constant $C_2 > 0$, which finishes the proof. \square

3.2. Confined subgroups. We recall the following definition.

Definition 3.3. Let G be a group. A subgroup H of G is **confined** if there exists a finite subset $P \subset G \setminus \{1\}$ such that $H^g \cap P \neq \emptyset$ for all $g \in G$.

Equivalently, a subgroup H is confined if the closure of the set of conjugates of H in the space $\text{Sub}(G)$ of subgroups of G , does not contain the trivial subgroup. The following simple lemma explains the usefulness of the notion of confined subgroups for the study of growth of actions. See [LBMB22a, Lemma 1.8] for a proof.

Lemma 3.4. *Let G be a finitely generated group, and X a G -set. Let*

$$\mathcal{S}(X) = \overline{\{G_x : x \in X\}} \subseteq \text{Sub}(G).$$

Then for every $H \in \mathcal{S}(X)$, we have $\text{vol}_{G,G/H}(n) \leq \text{vol}_{G,X}(n)$. In particular if some G_x is not confined then $\text{vol}_{G,X}(n) \simeq \text{vol}_G(n)$.

We will invoke the following, which is a particular case of Proposition 1.6 in [LBMB22a].

Lemma 3.5. *Let G be a finitely-generated group, and let $H \leq K$ be subgroups of G . Then $\text{vol}_{G,G/K} \leq \text{vol}_{G,G/H}$, and if in addition H has finite index in K , then $\text{vol}_{G,G/H} \simeq \text{vol}_{G,G/K}$.*

3.3. Houghton groups. If Ω is a set, $\text{Sym}(\Omega)$ is the group of all self-bijections of Ω . We denote by $\text{Alt}_f(\Omega)$ the group of alternating finitely supported permutations of Ω . For $H \leq \text{Sym}(\Omega)$, write $\Omega_{H,f}$ for the union of finite orbits under H , and $\Omega_{H,\infty}$ for the union of infinite ones. We say that a partition is H -invariant if every element of H sends a block of the partition to a (possibly different) block of the partition.

The following is a special case of [LBMB22b, Proposition 3.7].

Proposition 3.6. *Let Ω be a countable set, and let $G \leq \text{Sym}(\Omega)$ be any subgroup containing $\text{Alt}_f(\Omega)$. Then a subgroup $H \leq G$ is confined if and only if $\Omega_{H,f}$ is finite and there exists an H -invariant partition $\Omega_{H,\infty} = \Omega_1 \cup \dots \cup \Omega_k$ into infinite subsets such that H contains $\text{Alt}_f(\Omega_1) \times \dots \times \text{Alt}_f(\Omega_k)$.*

For $r \geq 2$, let Ξ_r be the graph obtained by gluing r infinite rays ℓ_1, \dots, ℓ_r at their origin. The Houghton group H_r is the group of all permutations g of the vertex set of Ξ_r , such that there exists finite subset $E, F \subset \Xi_r$ such that for every i , $g|_{\ell_i \setminus E}$ is an isometry onto $\ell_i \setminus F$. The group H_r is finitely generated for $r \geq 2$. Since the action of H_r on Ξ_r is by permutations of bounded displacements, it satisfies $\text{vol}_{H_r, \Xi_r}(n) \simeq n$.

In what follows, we fix $r \geq 2$. Let us denote Ω the vertex set of Ξ_r , and by $A := \text{Alt}_f(\Omega)$. Recall that A is simple and contained in every non-trivial normal subgroup of $\text{Sym}(\Omega)$. In particular A is a normal subgroup of H_r (and the quotient H_r/A can easily be seen to be isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}^{r-1}$).

Proposition 3.7. *Let $G := H_r$ be the Houghton group, with $r \geq 2$. Let X be a faithful G -set such that $\text{vol}_{G,X}(n) \not\asymp n^2$. Let $X = \sqcup_{i \in I} X_i$ be the decomposition of X into G -orbits. Then for every i either the action of G on X_i is not faithful (equivalently, the group A acts trivially), or it is conjugate to the diagonal action on $\Omega \times Y$, where Y is a finite G -set whose cardinality is uniformly bounded (that is, the upper bound that does not depend on the choice of the orbit X_i).*

Proof. We fix a word metric $\|\cdot\|$ on G , associated to some symmetric generating set. Let X be a G -set as in the statement and K be the stabiliser of a point $x_0 \in X$ such that A acts non-trivially on the orbit X_i of x . The group G has exponential growth, so by Lemma 3.4, K must be confined. Thus, we can apply Proposition 3.6.

Let us first show that $|\Omega_{K,f}| \leq 1$. Suppose by contradiction that this is not the case. Let $K_0 \leq K$ be the pointwise stabilizer of $\Omega_{K,f}$, which has finite index. Then $\text{vol}_{G,G/K_0}(n) \simeq \text{vol}_{G,X_i}(n)$ by Lemma 3.5. Choose two distinct points $x_1, x_2 \in \Omega_{K,f}$ and let $K_1 \geq K_0$ be

their pointwise stabiliser. Since $\text{vol}_{G, X_i}(n) \geq \text{vol}_{G, G/K_1}(n)$, a contradiction will follow if we show that $\text{vol}_{G, G/K_1}(n) \geq n^2$. Since the action of G on Ω is highly transitive (that is, transitive on n -tuples of distinct points), upon replacing K_1 by some G -conjugate we suppose that x_i is the point at position 1 on ℓ_i for $i = 1, 2$ (with the convention that the common origin is at position 0).

Now for $i = 1, 2$, let σ_i be the transposition that swaps the first and second positions of ℓ_i . Let also $t_i \in G$ be any element such that $t_i|_{\ell_i}$ shifts ℓ_i inside itself by 1. For example, these can be taken to be t, t^{-1} where t is the shift along a \mathbb{Z} -isomorphic ray coming from juxtaposing the rays ℓ_1, ℓ_2 . For $i = 1, 2$ and $n \geq 1$ consider the element

$$\gamma_{i,n} = (t_i^n \sigma_i t_i^{-n})(t_i^{n-1} \sigma_i t_i^{-n+1}) \cdots \sigma_i = t_i^n \sigma_i (t_i^{-1} \sigma_i)^n.$$

Note that $\|\gamma_{i,n}\| \leq Cn$, with $C = \max\{\|\sigma_i\|, \|t_i\| \mid i = 1, 2\}$. On the other hand each $\gamma_{i,n}$ is a permutation with finite support contained in ℓ_i , and $\gamma_{i,n}(x_i)$ is the point at position $n + 1$ on ℓ_i . Hence applying products of the form $\gamma_{1,m_1} \gamma_{2,m_2}$ to the pair (x_1, x_2) , with $0 \leq m_1, m_2 \leq n$ shows that $\text{vol}_{G, G/K_1}(n) \geq n^2$. This is a contradiction and proves the claim that $|\Omega_{K,f}| \leq 1$.

We now consider the partition $\Omega_{K,\infty} = \Omega_1 \cup \cdots \cup \Omega_k$ and claim that $k = 1$. Suppose that $k \geq 2$. By a similar reasoning as above (passing first to a finite index subgroup of K , and then to an overgroup), we have $\text{vol}_{G, X}(n) \geq \text{vol}_{G, G/K_1}(n)$, where K_1 is the subgroup of G of elements that preserve both Ω_1 and Ω_2 , so a contradiction will follow if we show that $\text{vol}_{G, G/K_1}(n) \geq n^2$. Fix n large enough. Again by high transitivity and using the fact that Ω_1, Ω_2 are both infinite, we can find $g \in G$ such that the point at position 1 on ℓ_1 is in $g(\Omega_1)$, and all points on ℓ_1 at positions between 2 and n are in $g(\Omega_2)$, while the symmetric statement holds for ℓ_2 . Now applying to $g(\Omega_1)$ and $g(\Omega_2)$ the same the elements $h = \gamma_{1,m_1} \gamma_{2,m_2}$ defined above with $1 \leq m_i \leq n$, we obtain n^2 different pairs $(hg(\Omega_1), hg(\Omega_2))$ with $\|h\| \leq 2Cn$. This shows that the ball of radius $2Cn$ around gK_1 has cardinality at least n^2 . Hence $k = 1$.

It follows that K contains $\text{Alt}(\Omega_{K,\infty})$. This also shows that $\Omega_{K,f} \neq \emptyset$, since otherwise K would contain the normal subgroup A , contradicting that the action on X_I is faithful. Hence $K \cap A$ contains a point stabiliser for the action of A on Ω , and hence is equal to it, as the latter is a maximal subgroup of A . Since K was an arbitrary point stabiliser, we deduce that the action of A on X_i is conjugate on each of its orbits to the standard action of A on Ω . For each A -orbit $\mathcal{O} \subset X_i$ we denote by $j_{\mathcal{O}}: \mathcal{O} \rightarrow \Omega$ the unique A -equivariant bijection (its uniqueness follows from the fact that the only A -equivariant permutation of Ω is the identity).

Let $Y = X_i/A$, the space of A -orbits, on which G acts since A is normal. For a point $x \in A$, we denote by $\mathcal{O}(x)$ its A -orbit. Consider the map $\iota: X_i \rightarrow \Omega \times Y$ given by $\iota(x) = (j_{\mathcal{O}(x)}(x), \mathcal{O}(x))$. This map is equivariant (this is obvious for the second component; for the first, it follows from the observation that the map $g^{-1} \circ j_{\mathcal{O}(gx)} \circ g$ is an A -equivariant bijection from $\mathcal{O}(x)$ to Ω , and thus it is equal to $j_{\mathcal{O}(x)}$). The map ι is also clearly injective. To check that it is surjective it is enough to observe that the diagonal G -action on $\Omega \times Y$

is transitive: this is true because the G -action on Y is transitive, and the group A acts trivially on Y and transitively on each fiber $\Omega \times \{y\}$.

To conclude we need to argue that Y must have uniformly bounded cardinality. Fix n and suppose that the diameter of Y is at least n . Then we can find a point $y \in Y$ and elements s_1, \dots, s_n in the generating set of G such that, if we set $g_m = s_m \cdots s_1$, then the points $y, g_1 y, \dots, g_n y$ are pairwise distinct (the point y and sequence (s_i) can be found by looking at any path in the Schreier graph of Y of length at least n). Next consider again the elements $\gamma_{i,n}$ above. Note that for every n , the element $\delta_n = \gamma_{1,n} \gamma_{2,n}$ belongs to A and thus acts trivially on Y , and coincides with $\gamma_{1,n}$ in restriction to $\ell_1 \subset \Omega$. Let $x \in \ell_1$ be at position 1, and consider the point $(x, y) \in \Omega \times Y$. Then the n^2 points $g_{m_2} \delta_{m_1}(x, y) = (g_{m_2} \gamma_{1, m_1} x, g_{m_2} y)$ for $m_1, m_2 \leq n$ are pairwise distinct, showing that there are at least n^2 in a ball of radius $O(n^2)$. This can't be true for n arbitrarily large, showing that the cardinality of Y must be uniformly bounded. \square

Theorem 3.8. *For every $r \geq 2$ and every non-trivial finitely generated group K , the group $H_r * K$ has a Schreier growth gap n^2 .*

Proof. By Proposition 3.2, it is enough to show that $G = H_r$ satisfies the sparse support condition at scale (n, n^2) . This is an immediate consequence of Proposition 3.7. Indeed fix any non-trivial element $g \in A$. Then for every faithful G set X such that $\text{vol}_{G,X}(n) \not\asymp n^2$, the element g has finite support in each G -orbit, so one can find constants $C, D > 0$ such that g satisfies conditions (C1)–(C3) for every $R > 0$, and condition (C4) is automatic, because every infinite connected graph has at least linear growth. \square

3.4. Topological full groups. Throughout this section we let Z be a Cantor space, and $\mathcal{G} \leq \text{Homeo}(Z)$ be a subgroup of its group of homeomorphisms. Recall that the topological full group of \mathcal{G} is the group $F(\mathcal{G})$ of all homeomorphisms h of Z such that for every $z \in Z$, there exist a clopen neighbourhood U of z and $g \in \mathcal{G}$ such that $h|_U = g|_U$. When \mathcal{G} is a cyclic group generated by a single homeomorphism φ , we write $F(\varphi)$ instead of $F(\mathcal{G})$.

We say that an element $g \in F(\mathcal{G})$ is a 3-cycle if $g^3 = 1$ and the set $\text{supp}(g) := \{z : g(z) \neq z\}$ is clopen and can be partitioned into 3 distinct clopen subsets U_1, U_2, U_3 such that $g(U_i) = U_{i+1}$ (with $i \bmod 3$). Following Nekrashevych [Nek19], the subgroup of $F(\mathcal{G})$ generated by 3-cycles is called the alternating full group of \mathcal{G} , and is denoted $A(\mathcal{G})$. It is shown in [Nek19] that if the action of \mathcal{G} on Z is minimal (that is, all its orbits are dense), then the group $A(\mathcal{G})$ is simple and contained in every non-trivial normal subgroup of $F(\mathcal{G})$. It is also shown in [Nek19] that if \mathcal{G} is finitely generated and its action on Z is expansive and all its orbits have cardinality at least 5, then $A(\mathcal{G})$ is finitely generated. Recall that an action of a finitely generated group \mathcal{G} on the Cantor space Z is expansive if and only if it is conjugate to a subshift (a \mathcal{G} -invariant subset of $\{1, \dots, d\}^{\mathcal{G}}$ for some $d \geq 2$). In many cases, results of Matui show that $A(\mathcal{G})$ coincides with the commutator subgroup of $F(\mathcal{G})$ (see his survey [Mat17]).

It is well-known (and not difficult to see) that if the group \mathcal{G} is finitely generated, then for every finitely generated subgroup G of $F(\mathcal{G})$, the identity map on Z defines a Lipschitz map between the graphs of the actions $\Gamma(G, Z) \rightarrow \Gamma(\mathcal{G}, Z)$. Since the groups $A(\mathcal{G})$ and \mathcal{G} have the same topological full groups, it follows that if the action of \mathcal{G} on Z is expansive and

has no orbits of cardinality less than 5, then the graph $\Gamma(A(\mathcal{G}), Z)$ is bi-Lipschitz equivalent to $\Gamma(\mathcal{G}, Z)$. In particular $\text{vol}_{A(\mathcal{G}), Z}(n) \simeq \text{vol}_{\mathcal{G}, Z}(n)$.

For a group G acting on Z by homeomorphisms, and for a finite set $Q = \{z_1, \dots, z_r\} \in Z$ we denote by G_Q the setwise stabiliser of Q , and, and by G_Q^0 its subgroup consisting of elements fix pointwise some neighbourhood of every $z \in Q$. The following is proven in [MB18].

Theorem 3.9 ([MB18]). *Let $\mathcal{G} \leq \text{Homeo}(Z)$ be a group acting minimally on Z , and let $G = A(\mathcal{G})$. Then a subgroup H of G is confined if and only if there exists a finite set $Q \subset Z$ such that $G_Q^0 \leq H \leq G_Q$ (the set Q is moreover unique).*

Corollary 3.10. *Retain the assumptions of Theorem 3.9, and let $\alpha(n) = \text{vol}_{\mathcal{G}, Z}(n)$. Let X be a faithful G -set such that $\text{vol}_{G, X}(n) \not\asymp \alpha(n)^2$, and let H be the stabiliser of a point in X . Then there exists $z \in Z$ such that $G_z^0 \leq H \leq G_z$.*

Proof. The assumption implies that H must be confined (Lemma 3.4). Let $Q \subset Z$ be the finite subset given by Theorem 3.9. Enumerate the points of Q as z_1, \dots, z_r and choose pairwise disjoint clopen neighbourhoods U_i of z_i . For each i we choose a finitely generated subgroup $G_i \leq G$, supported on U_i , such that the inclusion map $U_i \rightarrow Z$ defines a quasi-isometry of graphs $\Gamma(G_i, U_i) \rightarrow \Gamma(G, Z)$ (one can choose G_i to be the alternating full group of the restriction of the groupoid of germs of \mathcal{G} to U_i ; its finite generation follows from [Nek19, Proposition 5.4] and the quasi-isometry of the graphs from [Nek15, Cor 2.3.4]). It follows that the action of $G_1 \times \dots \times G_r$ on the orbit of Q has growth bounded below by $\alpha(n)^r$, from which it follows that $\text{vol}_{G, G/H}(n) \geq \text{vol}_{G, G/G_Q}(n) \geq \alpha(n)^r$, which contradicts the assumption unless $r = 1$. \square

Recall that for any subgroup $G \leq \text{Homeo}(Z)$, the maps $Z \rightarrow \text{Sub}(G)$ defined by $z \rightarrow G_z$ and $z \rightarrow G_z^0$ are respectively upper and lower semi-continuous. When G is countable, semi-continuity implies that $G_z = G_\delta$ for z in a dense G_δ subset.

Theorem 3.11. *Assume that \mathcal{G} is finitely generated and its action on Z is minimal and expansive, and let $\alpha(n) = \text{vol}_{\mathcal{G}, Z}(n)$. Then for every non-trivial finitely generated group H , the group $A(\mathcal{G}) * H$ has a Schreier growth gap $n\alpha(n)$.*

Proof. We shall show that G satisfies the sparse support condition at scale $(\alpha(n), \alpha(n)^2)$. As usual we fix once and for all a finite symmetric generating set S for G , which is used to define all Schreier graphs and distances below. Let X be a faithful G -set with $\text{vol}_{G, X}(n) \not\asymp \alpha(n)^2$. We shall show that it satisfies the conclusion of Definition 3.1 with constants $D = C = 2$. To this end, fix $R > 2$.

By [LBMB23, Lemma 6.2], for every G -orbit \mathcal{O} in Z , we have $\text{vol}_{G, \mathcal{O}}(n) = \alpha(n)$. Hence we can find a point $z_0 \in Z$ satisfying the generic condition that $G_{z_0}^0 = G_{z_0}$ and such that $|B_Z(z_0, R)| = \alpha(R)$. Choose $s_1, s_2 \in S$ such that $z_0, s_1(z_0), s_2 s_1(z_0)$ are distinct. Let (V_n) be a system of clopen neighbourhoods of z_0 , and such that $V_n, s_1(V_n), s_2 s_1(V_n)$ are disjoint for every n . Define accordingly the sequence (h_n) of 3-cycles given by

$$h_n(z) = \begin{cases} s_1(z) & z \in V_n \\ s_2(z) & z \in s_1(V_n) \\ s_1^{-1}s_2^{-1}(z) & z \in s_1s_1(V_n) \\ z & \text{else.} \end{cases}$$

Observe that $d_{G,Z}(z, h_n(z)) \leq 2$ for every $z \in Z$, by construction. The supports $\text{supp}(h_n)$ decreases to $\{z_0, h_n(z_0), h_n^2(z_0)\}$ as $n \rightarrow \infty$. It follows that for n is sufficiently large, we have $B_Z(z_0, R) \cap \text{supp}_Z(h_n) = \{z_0, s_1(z_0), s_2s_1(z_0)\}$. This shows that for n large, the element $g = h_n$ satisfies the conditions in Definition 3.1 for the action on Z .

Now consider the action on X and pick a point $x \in X$, and denote by $\phi(x) \in Z$ the unique point such that $G_{\phi(x)}^0 \leq G_x \leq G_{\phi(x)}$, given by Corollary 3.10. Note that $\text{supp}_X(h_n) \subset \phi^{-1}(V_n \cup s_1(V_n) \cup s_2s_1(V_n))$, since $h_n \in G_{\phi(x)}^0$ for $\phi(x) \notin V_n \cup s_1(V_n) \cup s_2s_1(V_n)$. However since the point z_0 satisfies $G_{z_0}^0 = G_{z_0}$ (and so do $s_1(z_0), s_2s_1(z_0)$) then it follows from semi-continuity and the inclusions $G_{\phi(x)}^0 \leq G_x \leq G_{\phi(x)}$ that for n large enough every x such that $\phi(x) \in V_n$, then the ball $B_X(x, R)$ will be isomorphic to the corresponding ball $B_Z(z_0, R)$, and similarly if $\phi(x) \in s_1(V_n), s_2s_1(V_n)$, with z_0 replaced by the corresponding image. Therefore the element $g = h_n$ satisfies all items in Definition 3.1 for the action on X as well. \square

REFERENCES

- [Ago13] Ian Agol, *The virtual Haken conjecture*, Doc. Math. **18** (2013), 1045–1087, With an appendix by Agol, Daniel Groves, and Jason Manning. MR 3104553
- [AS07] Goulnara Arzhantseva and Zoran Sunić, *Constructions of elements in the closure of grigorchuk group*, Geom. Dedicata **124** (2007), 5–26, Appendix to “The true prosoluble completion of a group: Examples and open problems” by G. Arzhantseva, P. de la Harpe, D. Kahrobaei, and Z. Šunić. MR 2318535
- [BG00] L. Bartholdi and R. I. Grigorchuk, *On the spectrum of Hecke type operators related to some fractal groups*, Tr. Mat. Inst. Steklova **231** (2000), no. Din. Sist., Avtom. i Beskon. Gruppy, 5–45.
- [Gri84] R. I. Grigorchuk, *Degrees of growth of finitely generated groups and the theory of invariant means*, Izv. Akad. Nauk SSSR Ser. Mat. **48** (1984), no. 5, 939–985. MR 764305
- [HW08] Frédéric Haglund and Daniel T. Wise, *Special cube complexes*, Geom. Funct. Anal. **17** (2008), no. 5, 1551–1620. MR 2377497
- [LBMB22a] A. Le Boudec and N. Matte Bon, *Growth of actions of solvable groups*, 2022, arXiv preprint.
- [LBMB22b] Adrien Le Boudec and Nicolás Matte Bon, *Confined subgroups and high transitivity*, Ann. H. Lebesgue **5** (2022), 491–522. MR 4443296
- [LBMB23] ———, *A commutator lemma for confined subgroups and applications to groups acting on rooted trees*, Trans. Amer. Math. Soc. **376** (2023), no. 10, 7187–7233. MR 4636688
- [Mat06] Hiroki Matui, *Some remarks on topological full groups of Cantor minimal systems*, Internat. J. Math. **17** (2006), no. 2, 231–251.
- [Mat17] ———, *Topological full groups of étale groupoids*, Operator algebras and applications—the Abel Symposium 2015, Abel Symp., vol. 12, Springer, [Cham], 2017, pp. 203–230. MR 3837599
- [MB18] N. Matte Bon, *Rigidity properties of topological full groups of pseudogroups over the cantor set*.
- [Nek15] Volodymyr V. Nekrashevych, *Hyperbolic groupoids and duality*, Mem. Amer. Math. Soc. **237** (2015), no. 1122, v+105. MR 3399889

- [Nek18] Volodymyr Nekrashevych, *Palindromic subshifts and simple periodic groups of intermediate growth*, Ann. of Math. (2) **187** (2018), no. 3, 667–719.
- [Nek19] V. Nekrashevych, *Simple groups of dynamical origin*, Ergodic Theory Dynam. Systems **39** (2019), no. 3, 707–732.
- [Neu37] B. H. Neumann, *Some remarks on infinite groups*, J. London Math. Soc. **12** (1937), 120–127.
- [Sal21] V. Salo, *Graph and wreath products in topological full groups of full shifts*, arXiv:2103.06663.
- [Sch27] O. Schreier, *Die Untergruppen der freien Gruppen*, Abh. Math. Sem. Univ. Hamburg **5** (1927), no. 1, 161–183.

CNRS, UMPA - ENS LYON, 46 ALLÉE D'ITALIE, 69364 LYON, FRANCE
E-mail address: `adrien.le-boudec@ens-lyon.fr`

CNRS, INSTITUT CAMILLE JORDAN (ICJ, UMR CNRS 5208), UNIVERSITÉ CLAUDE BERNARD LYON
1, 43 BLVD. DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE, FRANCE
E-mail address: `mattebon@math.univ-lyon1.fr`

UNIVERSITY OF TURKU, DEPARTMENT OF MATHEMATICS AND STATISTICS, FINLAND.
E-mail address: `vosalo@utu.fi`